

YENALIYEV, F.S.

Modification of Hess' operation in ptosis of the eyelids. Yest. oft. 73 no. 1:35-36 Ja-F '60. (MIRA 14:1) (EYELIDS—SURGERY)



APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962630010-3"



PETRASHEN', G.I.; YEMAL'SKIY, V.A.

Some interference phenomena in media containing thin horizontalparallel layers. Part 1. Isv.AN SSSR. Ser.geofix. no.9:1009-1020
s '56.


1. Akademiya nauk SSSR, Leningradskoye otdeleniye Matematicheskogo
instituta imeni V.A. Steklova.
(Seismic waves)


PETRASHEN', G.I.; TENAL'SKIY, V.A.

Some interference phenomena in media containing thin horizontal parallel layers. Part 2. Isv.AN SSSR.Ser.geofiz. no.10:1129-1144 (MIRA 10:1)

1. Akademiya nauk SSSR Leningradskoye otdeleniye Matematicheskogo instituta imeni V.A. Steklova.

(Seismic waves)






PETRASHEN', G.I.; YENAL'SKIY, V.A.

Some interference phenomena in media containing thin horizontalparallel layers. Fart 3. Izv.AN SSSR.Ser.geofiz. no.11:1241-1251 N 156. (MIRA 10:1)

indextrink intrinsial dancastrican nondextendiminations and an armanism of the contribution of the contrib

1. Akademiya nauk SSSR. Leningradskoye otdeleniye Matematicheskogo instituta imeni V.A. Steklova.

(Seismic waves)



APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962630010-3"

ACC NR: AP7004629

SOURCE CODE: UR/0288/66/000/003/0003/0012

AUTHOR: Yenal'sky, V. A.

ORG: none

TITLE: Numerical solution of the dispersion equation for beam-excited plasma

oscillations

SOURCE: AN SSSR. Sibirskoye otdeleniye. Izvestiya. Seriya tekhnicheskikh nauk, no. 3,

1966, 3-12

TOPIC TAGS: plasma beam interaction, plasma oscillation, plasma wave propagation,

dispersion equation, numerical solution, probability integral

ABSTRACT: A computer program is developed for solving the dispersion equation satisfied by the complex frequency $Z \sim \Omega + i\Gamma$ of longitudinal oscillations in the plasma, for the case in which thermal motion in the plasma and electron beam exhibits Maxwellian distribution. As this equation involves the probability integral of complex argument w(z), the program deals with 1) the computation of w(z) and its first two derivatives, and 2) the solution of the transcendental equation itself. Since classical w(z) programs for the construction of tables are not adequate for all values of z, an additional program is presented which uses a system of analytic series expansions which are given in detail. Programs for the identification of the roots and their precise computation are also given. Typical examples of numerical results

Card 1/2

UDC: 533.9.12

| follow in | the form of tables and curves giving the frequency Ω and the Γ as functions of the velocity ν_0 of the electron beam for pa | rticular |
|-------------------------|--|----------|
| Values of V. S. Imsh | the various physical parameters. The author thanks N. N. Yan mennik for formulating the problem and for their interest in t Orig. art. has: 2 figures and 1 table. | onko and |
| | 20/ SUBM DATE: none/ ORIG REF: 006/ OTH REF: 001/ | (/2) |
| | | |
| | | : |
| | | |
| | | |
| • | | |
| | | |
| | | |
| | | |
| Card 2/2 | | |

ACC NR: AT7004276

SUURCE CODE: UR/2517/66/074/000/0093/0106

AUTHOR: Yenal'skiy, V. A.

ORG: none

TITLE: On the motion of particles in an electromagnetic field

SOURCE: AN SSSR. Matematicheskiy institut. Trudy, v. 74, 1966. Raznostnyye metody resheniya zadach matematicheskoy fiziki (Difference methods for solving problems in mathematical physics), pt. 1, 93-106

TOPIC TAGS: electromagnetic field, particle motion, elliptic differential equation, iteration, numeric solution

ABSTRACT: The collisionless motion of particles of mass m and charge e in an electromagnetic field in free space is considered. Maxwell's equations with the Lorentz gauge and the equation of motion are reduced to the form (in cylindrical coordinates)

Card 1/2

ACC NRI AT7004276

where

 $cl = l', \quad \alpha = \frac{e}{mc^2}, \quad \frac{df}{dl'} = f.$

Here the scalar potential

$$\Phi = \overline{\Phi}_{\bullet} + \Psi_{\bullet}$$

has been separated into a high frequency component and a low frequency component $. \overline{\Phi}_{\bullet} = \Phi_{\bullet} \cos (\omega t + \theta).$

The motion of particles is considered in a cylindrical region of radius R and length L on whose surface & are given the boundary conditions

$$\Phi (r,0) = \Phi (r,L) = 0,$$

$$\Phi\left(R,z\right)=f\left(z\right),$$

$$\Psi|_{\Sigma}=\underline{0},$$

$$\Psi|_{\Sigma} = \underline{0},$$

$$A_{\diamond}(r, 0) = A_{\diamond}(r, L) = 0,$$

$$\sqrt{\frac{1}{r}} \frac{\partial}{\partial r} r \Lambda_{\bullet}|_{(R,x)} = H_{z}(z),$$

where f and Hz are given functions. The methods for numerical solution of the problem are discussed in detail, and the conditions for convergence of the iteration process are described. Some numerical results for several specific examples are presented and in one case are compared with the known analytic solution. The author thanks N. N. Yanenko, V. A. Teplyakov, and V. S. Imshennik for discussions, advice, and interest in the problem. A. A. Nikiforova helped write the computer programs. Orig. art. has: 124 equations, 3 figures, and 1 table.

Card 2/2 SUB CODE: 20/ SUBM DATE: none/ ORIG REF: 007/ OTH REF: 003

USSR/Thermodynamics - Thermochemistry. Equilibria.

B-8

Physical-Chemical Analysis. Phase Transitions.

Referat Zhur - Khimiya, No 6, 1957, 18530 Abs Jour

: A.I. Yurzhenko, V.D. Yenal'yev. Author

: Lvov University. Inst : Study of Distribution of Isppropylbenzene Hydroperoxide

Title between Styrene and Water Phases.

: Nauk. zap. L'vivs'k. un-tu, 1955, 34, 45-50 Orig Pub

The distribution of isopropylbenzene hydroperoxide (I) Abstract

between the styrene and water phases at 200, 350 and 430 was studied. The distribution factor (K) describing the ratio of molar parts of I in the water phase and in styrema decreases with the temperature rise from 0.0639 at 20° to 0.0314 at 43°. Addition of small amounts of NaOH (0.001 to 0.025 n.) causes a decrease of K due to salting out, but at the increase of NaOH concentration to 0.1 n., K rises due to the formation of a I salt soluble in water. Addition of Na₂CO₃ and K₂SO₄ causes salting

- 209 - out of I and a decrease of K. Card 1/1

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962630010-3


THREFIELD A.I.: TENAL TEV, V.D.


Interaction between organic hydroperoxides and ferrous salts.

Interaction between organic hydroperoxides and ferrous salts.

Dop. ta pov. L'viv. un. no.7 pt.;:195-197 '57. (MIRA 11:2)

(Chemical reaction, Rate of;
(Hydroxides) (Iron salts)





YURZHENKO, A.I. [IUrshenko, O.I.]; YENAL'TEV, V.D. [IBnal'iev, V.D.]

Investigation of the reactions of organic hydroperoxides with salts of ferrous oxide. Mauk.rap.L'viv.un 46:7-12 '58.

(Hydroperoxides) (Iron salts)

(Hydroperoxides)

KUCHER, R.V.; YENALIYEV, V.D. [IEnal iev, V.D.]; YURZHENKO, A.I., [IUrzhenko, O.I.], Kovbuz, M.O.

Effect of the molecular weight of tertiary hydrocarbons on their oxidizability in the liquid phase and in emulsions. Nauk. zap.L'viv.un. 46:13-16 '58. (MIRA 12:7) (Hydrocarbons) (Oxidation)

YENAL'YEV, V.D. [IEnal'iev, V.D.]; YUNZHENKO, A.I. [IUrzhenko, O.I.]

Effect of the relationship of phase to the kinetics of redox polynerization in emilsions. Kauk.zap.L'viv.un. 46:21-2; '58. (MIRA 12:7)

(Polymerization)

YENAL'TEV, V.D. [IEnal'iev, V.D.]; YURZHENKO, A.I. [IUrzhenko, O.I.]

Effect of the concentration of the initiating system on the kinetics of redox polymerization in emulsions. Mauk.zap.L'viv.un 46: 26-33 '58. (Polymerization)

(Polymerization)

ON THE STATE OF A STAT

5(4) SOV/20-123-2-32/50 Yurzhenko, A. I., Ivanova, N. Ya., AUTHORS: Yenal yev, Y. D. The Participation of the Emulsifier in the Oxidation Reduction TITLE: Initiation of Emulsion Polymerization (Uchastiye emul'gatora v okislitel'no-vosstanovitel'nom initsiirovanii emul'sionnoy polimerizatsii) Doklady Akademii nauk SSSR, 1958, Vol 123, Nr 2, pp 324-326 PERIODICAL: (USSR) One of the most important factors influencing the kinetics of ABSTRACT: polymerization in emulsions is the nature of the emulsifying agent. The nature of the emulsifier used influences not only the velocity of the polymerization process but also the properties of the polymer formed. When investigating emulsion polymerization in the presence of various emulsifiers, the authors noticed several particularities in the development of the polymerization process in connection with the application of cetyl pyridine bromide. In this case the part of the emulsifier is played not only by a purely colloidochemical factor. Investigation was carried out by the dilatometric method in a dilatometer which prevents contact between the polymerization Card 1/4

 The Participation of the Emulsifier in the Oxidation SOV/20-123-2-32/50 Reduction Initiation of Emulsion Polymerization

system and air. In the case of all experiments, the ratio between the hydrocarbon- and the aqueous-phase was 1 : 9. The hydroperoxide of isopropyl benzene served as initiator, and styrene was used as monomer. Polymerization kinetics was investigated at various temperatures. In the course of one of the test series sodium carbonate was introduced into the aqueous phase. The results obtained by the experiments are shown in a diagram. Conditions otherwise being equal, polymerization develops much more rapidly than if other classes of emulsifiers are used. Cetyl pyridine bromide warrants sufficiently rapid polymerization also at low temperatures (4 and 18°), which is not the case with other emulsifiers. If sodium carbonate is present in the aqueous phase, polymerization velocity passes through a maximum at increased temperatures. In the course of experiments carried out without sodium carbonate, polymerization increases with rising temperature, in which case linear dependence is conserved up to a rather high degree of polymerization. An addition of sodium carbonate and an increase of temperature acts in the same direction (increase of polymerization velocity). The velocity

Card 2/4

The Participation of the Emulsifier in the Oxidation SOV/20-123-2-32/50 Reduction Initiation of Emulsion Polymerization

of the polymerization process is due to the velocity of initiation. The decay of isopropyl benzene hydroperoxide in an aqueous solution is considerably accelerated by the introduction of cetyl pyridine bromide also if Na₂CO₃ is lacking.

This decay is still more accelerated if cetyl pyridine bromide and sodium carbonate are present at the same time. Data concerning the kinetics of this decay at various conditions are given by a diagram. An increase of temperature increases the initial velocity of polymerization and reduces the final yield of the polymer. Also an addition of sodium carbonate produces the same effect. A comparison between these and other data makes it possible to draw the following conclusion: The surface-active emulsifier may play a double rôle in emulsion polymerization: Firstly, it may act as an ordinary emulsifier stabilizing the original emulsion of the monomer, and, secondly, the emulsifier may have the functions of a polymerization activator by causing an induced decay of the hydroperoxide. There are 4 figures and 7 references, 4 of which are Soviet.

Card 3/4

The Participation of the Emulsifier in the Oxidation \$07/20-123-2-32/50 Reduction Initiation of Emulsion Polymerization

ASSOCIATION: L'vovskiy gosudarstvennyy universitet im. Ivana Franko

(L'vov State University imeni Ivan Franko)

PRESENTED: July 3, 1958, by P. A. Rebinder, Academician

SUBMITTED: May 16, 1958

Card 4/4

Initiation of the emulsion oxidation of isogropylbenzend and 1, 1-diphenylethane by hydrogen peroxide, Sbor. nauch. rab. Inst. fiz.~corg. khim. AM BSSR no.8:126-131 '(60. (MIRA 14:3)) 1. L'vovskiy gosudarstvennyy universitet im. I. Franko. (Cumene) (Ethane) (Hydrogen peroxide)

s/076/61/035/010/010/015 28291 B106/B230

54300 also 1375

Kucher, R. V., Kaz'min, S. D., and Yenal'yev, V. D.

AUTHORS:

Initiation of emulsion oxidation of alkylated aromatic

hydrocarbons by hydrogen peroxide TITLE:

Zhurnal fizicheskoy khimii, v. 35, no. 10, 1961, 2322 - 2327

TEXT: The authors investigated the initiation by hydrogen peroxide in PERIODICAL emulsion oxidation of isopropyl benzene, 1,1 diphenyl ethane, and 1-phenyl-1-p-tolyl ethane in the liquid phase, this problem being of great practical interest in the synthesis of hydroperoxide compounds. Oxidation was conducted at 85°C in "air lift" type glass vessels in which the reaction mixture was agitated by air bubbling in through a porous glass partition. For the aqueous phase, a 0.1 N soda solution was used in all tests. The volume ratio of the hydrocarbon phase to the aqueous phase was 1:3. At regular intervals, samples were taken and the hydroperoxide content was determined iodometrically by potentiometric titration (Ref. 5: see below). Oxidation of the alkylated aromatic hydrocarbons referred to proceeds in emulsion systems by autocatalysis. The effect of hydrogen Card 1/4

CIA-RDP86-00513R001962630010-3 APPROVED FOR RELEASE: 03/15/2001

3/076/61/035/010/010/015 28291 B106/B230

Initiation of emulsion oxidation of ...

peroxide on the process is very specific, and depends not merely on the character of the hydrocarbon but also, in a high degree, on the instant of adding the hydrogen peroxide. Constant initiation by adding hydrogen peroxide at short intervals intensifies the oxidation of 1,1-diphenyl ethane, whereas it inhibits the oxidation of cumene. In all oxidation processes investigated, the following common rules could be observed: when hydrogen peroxide was added at the beginning of the process, reaction rate and hydroperoxide yield were practically not affected; when, however, hydrogen peroxide was added at the final stage of oxidation after maximum concentration of hydroperoxide was attained, a rapid rise in reaction rate and hydroperoxide yield took place anew. From observations made the following conclusions were drawn: Initiation by hydrogen peroxide did not simply cause a rise in the concentration of chain radicals as had been frequently assumed in publications. Apparently, radicals formed by decomposition of ${\rm H_2^{0}_2}$ were not sufficiently active to start new chains by reacting with the hydrocarbon. With progressing oxidation,

products accumulated in the system acting as inhibitors on oxidation. With such inhibitors initiator radicals may react, thus eliminating the inhibiting effect. For this reason, initiation effect increases with Card 2/4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962630010-3"

environ amenda de comparation de la comparation del comparation de la comparation de

28291

s/076/61/035/010/010/015 B106/B230

Initiation of emulsion oxidation of ...

progressing reaction. This hypothesis was confirmed by an experiment in which hydrogen peroxide was introduced into a reaction retarded by an inhibitor. For this purpose, the oxidation of cumene was inhibited by adding a small quantity (0.01 g-mole/liter) of phenol breaking down the oxidation chains according to reaction $C_6H_5OH + R^{\circ} \rightarrow C_6H_5O^{\circ} + RH$.

C6H50° radicals are of low activity, and recombine. Adding hydrogen peroxide eliminated the inhibition of the reaction, and caused a steep rise of the oxidation rate. When during the reaction, oxidation products combine with initiator radicals to form radicals similar to chain radicals in their activity, initiation results in increasing the total oxidation rate. In the reverse case, the consumption of components reacting with initiator radicals is accelerated and the total reaction rate decreases. Also in this case, the effect of a brief initiation at the final stage of oxidation may be favorable for the process. The effect of an initiator therefore depends on the reactivity of the components of the reaction mixture. There are 3 figures and 7 references: 5 Soviet and 2 non-Soviet. The two references to English-language publications read as follows: Ref. 5: V. Kokatnur, M. Jelling, J. Amer. Chem. Soc., 63, Card 3/4

APPROVED FOR RELEASE: 03/15/2001

2**5291** \$/076/61/035/010/010/015 B106/B230

Initiation of emulsion oxidation of ...

1432, 1941; J. W. Fordham, H. L. Williams, Canad. J. Chem., 27B, 913, 1954.

L'vovskiy universitet im. Iv. Franko (L'vov University imeni ASSOCIATION:

Iv. Franko)

March 3, 1960 SUBMITTED:

Card 4/4

81410

s/020/60/132/06/35/068 B004/B005

5.3200

Yenal'yev, V. D. Kucher, R. V., Kazimin, S. D.,

AUTHORS:

On the Possibility of Increasing the Yield in Hydroperoxide

TITLE:

by Initiating the Cumene Oxidation With Hydrogen Peroxide

PERIODICAL:

Doklady Akademii nauk SSSR, 1960, Vol. 132, No. 6,

pp. 1348-1351

TEXT: The authors discuss the process of initiation of a chain reaction on the basis of papers by N. M. Emanuel! (Ref. 1) and N. N. Semenov (Ref.2).

In previous papers by the authors (Refs. 3, 4) it was observed that in the case of initiation of oxidation of isopropyl benzene by means of H202 the effect depends on the point of time of adding the initiator (Fig. 1A). An addition at the beginning of oxidation effects neither acceleration of the reaction nor reduction of the induction period. Only if H202 is added at later points of time when the reaction becomes slower, it effects an acceleration so that the hydroperoxide yield rises

Card 1/3

81410

On the Possibility of Increasing the Yield in Hydroperoxide by Initiating the Cumens Oxidation With Hydrogen Peroxide S/020/60/132/06/35/068 B004/B005

from 40 to 80%. Hence, the authors conclude that the by-products developing during oxidation exert an inhibiting influence which is eliminated by H_2O_2 . They confirmed this conclusion by adding phenol as an inhibitor the effect of which was really eliminated by H_2O_2 (Fig. 1B). Equations are written down for the kinetics of the reaction $A \to B \to C$, with the product B undergoing degenerated branching, and C interrupting the reaction product B undergoing degenerated branching, and C interrupting the reaction of the product by H_2O_2 is also branching, H_2O_2 is added, the more intensive the reaction rate for C). The later the H_2O_2 is added, the more intensive its initiating effect. There are 2 figures and 6 references: 5 Soviet

ASSOCIATION: L'vovskiy gosudarstvennyy universitet im. Ivana Frankc (L'vov State University imeni Ivan Franko)

Card 2/3

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962630010-3

81410

On the Possibility of Increasing the Yield in Hydroperoxide by Initiating the Cumene Oxidation With Hydrogen Peroxide

S/020/60/132/06/35/068 B004/B005

PRESENTED:

February 11, 1960, by V. N. Kondrat'yev, Academician

SUBMITTED:

February 9, 1960

Card 3/3

YENAL'YEY, V.D. [IEnal'iev, V.D.]; ZAYTSEVA, V.V.; SADOVSKIY, Yu.S. [Sadovs'kyi, IU.S.]; BATOG, A.Ye. [Batoh, A.IE.]; SADOVSKAYA, T.M. [Sadovs'ka, T.M.]

Thermal stability and initiating activity of substituted benzoyl peroxide. Khim.prom. [Ukr.] no.1:17-20 Ja-Mr 164. (MIRA 17:3)

BATOR, A.Ye.; Shedriff, A.M.; Yenanyray, V.B.; Romersevick, L.L.

Synthesis of some tert-anylperacylates. Ukr. khin. stur. 33
(And 17:10)

no.9:954-955 'fd.

1. Ukrainskiy nauchno-issledovatel'skiy institut plastichoukikh
mass, Donetsk.

APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962630010-3"

HAZAROVA, Z.F.; BATOG, A.Ye.; YEHAL!YEV, V.D.; ROMANTSEVICH, M.K.

Gondensation of tertiary amyl hydroperoxide with some carbonyl compounds. Zhur. ob. khim. 34 no.7:2430-2432 [MIRA 17:8]

1. Ukrainskiy naucimo-issledovateliskiy institut plastmass, Donetsk.

BATOG, A.Ye.; TATARSKAYA, I.M.; ECCHAROVA, Yu.Ye.; YENAL'YEV, V.D.; REMANTSEVICH, M.K.

Synthesis of peroxide and hydroperoxide of tertiary butyl.
Ukr.khim.zhur. 31 no.2:207-208 165. (MIRA 18:4)

3. Ukrainskiy nauchno-issledovateliskiy institut plasticheskikh mass, Donetsk.

YENALIYEV, V.D.; KONDHATOVICH, A.A.; GENDRIKOV, E.P.; DEDOVETS, G.S.

Swelling of the copolymer of styrene with divinyl benzene.
Plast. massy no.8:5-6 165.

(MIRA 18:9)

APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962630010-3"

YENAL'YEV, V.D.; ZAYTSEVA, V.V.; SADOVSKIY, Yu.S.; SADOVSKAYA, T.N.; SOROKINA, A.N.

Kinetics of styrene polymerization in the presence of some tert-amyl peracylates. Ukr. khim. zhur. 31 no.8:834-838 '65. (MIRA 18:9)

1. Ukrainskiy nauchno-issledovatel'skiy institut plasticheskikh mass.

YENAL'TEV, V.B.; YAYTSEVA, V.V.; SAECUSKIT, Yu.S.; SAECUSKIY, T.H.;

Polymerization of styrene initiated by bifunctional peroxides.

Vysokom. soed. 7 no.2:275-279 F '65.

(MIRA 18:3)

1. Ukrainskiy nauchno-issledovatel'skiy institut plastichoskikh mass.

ACC NRI AP6017975 SOURCE CODE: UR/0413/66/000/010/0079/0079 INVENTORS: Yonal'yev, V. D.; Demidenko, A. G. ORG: none TITLE: A method for obtaining granular polymers. Class 39, No. 181807 Zannounced by Ukrainian Scientific Research Institute of Plastics (Ukrainskiy nauchnoissledovatel'skiy institut plasticheskikh mass) SOURCE: Izobretoniya, promyshlennyye obraztsy, tovarnyye znaki, no. 10, 1966, 79 TOPIC TAGS: polymer, polycondensation, plastic, formaldehyde, phenol, alumosilicate, ABSTRACT: This Author Certificate presents a method for obtaining granular polymers. The method involves suspensional polycondensation of one or several mixed polar substances that enter the polycondensation reaction and form oil-insoluble products, such as phenolsulfo acids and formaldehyde, in a nonpolar dispersing medium. To strengthon the stability of the emulsion, structuring substances are added to the dispersing medium. These substances possess hydrophylic-hydrophobic properties or are capable of assuming hydrophylic-hydrophobic properties due to an addition of hydrophobilizing or hydrophylizing addenda, for instance alumosilicates, silica gel or organic salts of heavy metals. SUB CODE: 11/ SUBM DATE: 14Jan63 Card 1/1 UDC: 678.6.034


Ü CHMIRI Bulgaria CAPSGORY 1959. No. 85658 ASO, JOUR. : RZKhim., No. : Yenchev, D.; Stancheva, P. ; Higher Agricultural Institute "V. Kolarov". AUERGR I. JT. : Chemical Characterization of Manganese Ore TIME from the "Poteda" Mine. : Nauchi tr. Vissh. selskostop. in-t " V. Kolaonic. Pub. rov" - Plovaiv, 1956 (1958), 5, 169-195 : Samples of ore were investigated in order to develop the technology of its industrial utilization. Determinations were made of variations of basic composition, proportions of oxides and compounds of kn of different valency, and a procedure is proposed for the production of potassium permangamate. -- G. Vorob'yev. GARD: 43

JURYAVTSEVA, A. V.


"Investigations of Spectra of Conversion Electrons and Spectra of Positrons of the Europium Fraction."

report submitted for All-Union Conf on Nuclear Spectroscopy, Tbilisi, 14-22 Feb 64.

OTYAI, IGU (Joint Inst Nuclear Res; Leningrad State Univ)



APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962630010-3"



APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962630010-3"

s/078/60/005/06/07/030 B004/B014

5.2600 AUTHOR:

TITLE:

Yenchev, D. G.

The Influence of Some Ammonium-, Potassium-, and Sodium Salts Upon the Decomposition Rate of Ammonium Nitrite in

the Reaction NHAC1 + NaNO2 = NHANO2 + NaCl

Zhurnal neorganicheskoy khimii, 1960, Vol. 5, No. 6, PERIODICAL:

pp. 1234 - 1240

TEXT: The author investigated the influence of NH_4 , K_7 , and Na salts of various anions, namely, CES, Br, NO_3 , C1, CH_3COO_9 and SO_4^{27} upon the decomposition of ammonium nitrite. He describes the experimental arrangement (Fig. 1), in which the temperature was rigorously kept constant, and the amount of nitrogen liberated was measured every 7 sec. Experimental results are graphically shown in Figs. 2-4 (abscissa - duration of the experiment, ordinate - liberated N₂ in cm³). The separation of N_2 in the reaction of ammonium chloride with sodium nitrite attains

Card 1/4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962630010-3"

The Influence of Some Ammonium., Potassium., \$\, 5/078/60/005/06/07/030 and Sodium Salts Upon the Decomposition Rate \$\, 8004/E014 of Ammonium Nitrite in the Reaction NH_4C1 + NaNO_2 \Rightarrow NH_4NO_2 + NaC1

a measurable intensity at 57°C, while the addition of ammonium salts yields measurable N₂ volumes already at 56°C. The anions of the ammonium salts speed up the reaction in the lyotropic series $\text{CNS}^- > \text{Br}^- > \text{NO}_3^- > \text{Cl}^- > \text{SO}_4^2 > \text{CH}_3\text{COO}. \text{ The accelerating effect of NH}_4 \text{ salts} \\ \text{is explained by the increase in concentration of the NH}_4^+ \text{ ion, the concentration being a factor of the equation according to which the decemposition of NH}_4^{\text{NO}}_2 \text{ is most likely to occur: } d(\text{N}_2)^4/\text{dt} = \text{K}_1 \left[\text{NH}_4^+\right] \left[\text{NO}_2^-\right] \left[\text{HNO}_2\right].$ The anions of potassium salts slow down the decomposition rate of NH}_4^{\text{NO}}_2 \right] in the same order. The anions of sodium salts have a similar, slightly more retarding action. With them, Cl and SO4 change their place in the order. The strongly inhibiting action of the acetate ion is explained by the low reactivity of HNO2 in its presence. With respect to the inhibition of the decomposition of NH}_4^{\text{NO}}_2 \right] the cations range in the following order:

Card 2/4

The Influence of Some Ammonium., Potassium., 8/078/60/005/06/07/030 and Sodium Salts Upon the Decomposition Rate B004/B014 of Ammonium Nitrite in the Reaction NH₄Cl + NaNO₂ = NH₄NO₂ + NaCl

NH_A < K⁺ < Na⁺. The influence of the ions upon the decomposition rate is explained by their hydration. Ions with a relatively low solvation heat like CNS, Br, NO₃, NH_A exhibit a greater accelerating or lesser inhibiting effect, respectively. The hydration is regarded as a substitution of ions for water molecules in the quasicrystalline structure of water. This reaction goes along with a heat effect, with thermal conductivity increasing with the ionic radius. Thus, the higher decomposition rate in the presence of bromine ions, as compared with chlorine ions, is explained by the larger radius of Br in aqueous media, whereby the quasicrystalline structure of water is weakened and the motion of heat is facilitated. The cations and anions investigated had an additive effect. The author mentions papers by K. P. Mishchenko (Ref. 8), M. D. Lagunov (Ref. 9), N. Ye. Khomutov (Ref. 10), A. F. Kapustinskiy and I. I. Ruzavin (Ref. 12). There are 3 figures and 12 references: 6 Soviet, 5 German, and 1 Austrian.

Card 3/4

The Influence of Some Ammonium-, Potassium-, S/078/60/05/06/07/030 and Sodium Salts Upon the Decomposition Rate B004/B014 of Ammonium Nitrite in the Reaction NH4C1 + NaNO2 = NH4NO2 + NaC1

ASSOCIATION: Vyushiy sel'skokhozyaystvennyy institut im. V. Kolarova,

Kafedra obshchey khimii, Plovdiv (Bolgariya) (Higher

Agricultural Institute imeni V.Kolarov, Chair or General

Chemistry, Plovdiv (Bulgaria))

January 13, 1959 SUBMITTED:

Card 4/4

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962630010-3

VENCH BULDARIA/Diseases of Farm Animals - Diseases Caused by Viruses R-3 and Rickettsiae.

Abs Jour

: Ref Zhur - Biol., No 14, 1958, 64656

Author

: Ivanov, Ks., Zhelev, Vl., Yenchev, St.

Inst

: Institute of Experimental Vaterinary Medicine of the Bulga-

rian Academy of Sciences.

Title

The Study of the Morphological Changes in Swine Plague in

Relation to the Diagnostic Criteria of this Discaso.

8. Changes in the Storach and Intestines.

Orig Pub

Izv. In-ta eksperim. vct. ncd. B"lgar. AN, 1956, 4, 213-

231.

Abstract

The authors analyze the results of the investigation of the gastrointestinal tract in 196 pigs sacrificed on the 4th-

6th day following experimental infection with plague, and in 19 swine which perished from spontaneous acute plague.

Card 1/2

- 18 -

BULGARIA/Discases of Farm Animals - Discases Caused by Viruses R-3
and Rickettsiaes

Abs Jour : Ref Zhur - Biol., No 14, 1958, 64656

They observe that in acute plague hyperemia, hemorrhages and lymphoidocytic infiltrates are encountered in the mucosa more often than it is usually accepted. Besides hemorrhagic diathesis, the plague virus of swine also causes the necrotic processes, especially in the glandular part of the stomach and in the ilcum, leading to the formation of the impregnated with fibrin, circumscribed, button-like, false membranes or sores. The authors do not consider as characteristic of the swine plague such changes in the large intestine as the inflammation of the terminal part of it, and the cosinophilia in its submucosa.

Card 2/2

USSR/General Biology - Genetics. Plants Genetics.

B.

Abs Jour

: Ref Zhur - Biol., No 21, 1958, 94692

Author

Title

: Yenchev, Yanko

Inst

: Adequacy of Changes in Heridity During Transformation of

Spring Forms in Winter.

Orig Pub : Agrobiologiya, 1957, No 5, 101-105

Abstract

: The author transformed spring forms of wheat and barley into winter by means of seeding before winter. According to his data, the crythrospermum variety was transformed into the lutescens; in the lutescens variety 1163 varieties of crythrospermum and hilturum appeared. -- S.Ta.

Krayevoy

Card 1/1

- 34 -

YENCHEV. Ya., kand.biolog.nauk; MOSKOV, I.; BOZOVA, L.

Developing Eritrospersum 341 spring wheat into frost resistant winter wheat. Dokl.Akad.sel'khoz. 24 no.6:10-11 '59. (MIRA 12:9)

1. Sel'skokhozyaystvennyy institut im. G.Dimitrova, Bolgariya, g.Sofiya. Predstavlena akademikom M.A.Ol'shanakim. (Wheat breeding)

TENCHEV, Ya., kend.bjol.nenk, MOSKOW, I.; BOZOVA, L. (Bolgariya)

Changes in anatomical and physiological properties of wheat when transformed from spring varieties into winter varieties. Agrebiotogiia no.6:266-872 N-D '60.

1. Sel'skokhozyaystvennyy institut imeni G.Dimitrova, g. Sofiya. (Wheat)

APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962630010-3"

YENCHEV, Ya.; MOSKOV, I.; BOZOVA, L.

Developing heritable winter hardiness in the Mutans-103 barley by controlled cultivation. Dokl. AN SSSR 135 no.6:1536-1538 D '60. (MRA 13:12)

1. Sel'skokhozyaystvennyy institut im. G. Dimitrova, Sofiya, Bolgariya. Predstavleno akadmikom T.D. Lysenko. (Barley) (Plants--Frost resistance) (Heredity)

YENCHEV, Ya. (Narodnaya Respublika Bolgarii); BOZOVA, L. (Narodnaya Respublika Bolgarii)

Physiological and biochemical changes in the process of transforming spring varieties into winter varieties. Agrobiologiia no.6:803-814 N-D '63. (MIRA 17:2)

1. Seliskokhozyaystvennyy institut imeni G. Dimitrova, Sofiya.

YFNCHEV, Ya.: BOZOVA, L.

Induced physiological and biochemical changes in the process of the transformation of spring into winter varieties of wheat. Agrobiologiia no.5:671-680 S-0 165.

(MIRA 18:9)


1. Vysshiy sel'skokhozyaystvennyy institut imeni G.Dimitrova, Sofiya, Bolgariya.

DANILOV, V.I.; YENCHEVICH, I.B.; NOVIKOV, D.L.; POLFEROV, E.A.; SAFONOV, A.N.; FEOKTISTOV, B.V.


[Calculation of the initial region of stable phase oscillations in a synchrocyclotrone] Raschet nachal'noi oblasti ustoichivykh kolebanii v sinkhrotsiklotrone. Dubna, Ob"edinennyi in-t iadernykh issl. 1963. 24 p. (MIRA 17:7)


DANILOV, V.I.; YENCHEVICH, I.B.; ZAMOLODCHIKOV, B.I.; MARCHENKO, B.N.; NOVIKOV, D.L.; POLFEROV, E.A.; ROZANOV, Ye.I.; SAVENYOV, A.L.; SAPONOV, A.N.


Increase in intensity of a proton beam in a six-meter synchro-cyclotron of the United Institute of Nuclear Research. Atom. energ. 16 no.1:9-11 Ja '64. (MIRA 17:2)





APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962630010-3"

















| ATTROVED TO RELEASE US/13/13/13/13/13/13/13/13/13/13/13/13/13/ |
|---|
| |
| L LLL-66 ENT(m)/EFA(w)-2/ENA(m)-2 IJP(c) DM -ACCESSION NR: AP5023773 UR/0089/65/019/003/0289/0292 -621.384.611 |
| AUTHOR: Danilov, V. I.; Yenchevich, I. B.; Zamolodchikov, B. I.; Polferov, E. A.; Rozanov, Ye. I.; Smirnov, V. I.; Testov, V. G. |
| TITLE: The increase in pulse duration of the 680 MEV OIYal synchrocyclotron particle beam |
| SOURCE: Atomnaya energiya, v. 19, no. 3, 1965, 289-292 TOPIC TAGS: synchrocyclotron, ion acceleration, ion accelerator, MEV accelerator |
| ABSTRACT: In synchrocyclotrons icas are accelerated in bunches, the shape and dimensions of which are determined by radial-phase and betatron oscillations. The present authors describe a method for pulse extension which was tested on the OIYaI synchrocyclo-authors describe a method for pulse extension which was tested on the OIYaI synchrocyclo-authors describe a method for pulse extension which was tested on the OIYaI synchrocyclo-authors describe a method for pulse extension. |
| $T = \int_{t_{M}}^{\infty} \frac{dr}{\hat{r}_{\theta}(t) + \hat{q}_{\theta}, y_{\theta}(t)}$ |
| Card 1/4 |
| |

L 4147-66 ACCESSION NR: AP5023773

where the speed of equilibrium orbit widening is given by

$$\dot{r}_{s} = \frac{r_{s}}{1-n} \cdot \frac{1}{E_{s}\beta_{s}^{2}} \cdot \frac{\omega_{s}}{2\pi} \epsilon_{0} V_{0} \sin \varphi_{s} = \frac{r_{s}}{1-n} \cdot \frac{1}{K_{s}\beta_{s}^{2}\omega_{s}} \cdot \frac{d\omega}{dt};$$

 $f_{B,M}$ (t) is velocity of displacement of the equilibrium orbit at the f_n azimuth caused by the excitation of the first harmonics of the magnetic field;


$$n = -\frac{r}{H} \cdot \frac{\partial H}{\partial r} : K = 1 + \frac{n}{1 - n} \cdot \frac{1}{\beta^3} : \beta = \frac{v}{\sigma} :$$

v, ω , E are velocity, rotational frequency, and total energy of the particle, respectively; eV_0 - maximum possible energy increment per turn; subscripts a characterize equilibrium values;

$$\delta r = q_0 + q_0$$
 with $q_{a.M} = 0$;
 $\delta r = q_0 + 2q_0$ with $r_a = 0$

Card 2/4

| -3.1.1 | | |
|---------|--|--------------|
| | | |
| | L 4147-66 | |
| 3/2.741 | ACCESSION NR: 1.5023773 | |
| | and \int_{C} is the maximum amplitude of radial betatron and radial-phase oscillations respectively. It is shown that the length of the pulse may be extended by increasing the interval of radial oscillation amplitudes and by decreasing the beam velocity along the radius (this can be achieved by increasing, in time, the forced radial oscillations for $r_{c} = 0$). A brief description of the design and operation of the necessary circuits is also | |
| | g = 0), A brief description of the design and 5 floures. | |
| | given. Orig. art. has: 9 formulas and 5 figures. | |
| | ASSOCIATION: None | |
| 34 | CUBMITTED: 06Feb65 ENCL: 01 SUB CODE: NP, MA | |
| | NO REF SOV: CO1 OTHER: 006 | |
| | | |
| - 100 | | |
| | Card 3/4 | |
| | | in Leave the |
| | | |




EWT (m) IJP(c) 07919-67 SOURCE CODE: UR/0120/66/000/003/0019/0022 ACC NR: AP6021991 AUTHOR: Danilov, V. I.; Yenchevich, I. B.; Rozanov, Ye. I.; Tomilina, T. M.; Shestov. A. V. ORG: Joint Nuclear Research Institute, Dubna (Ob"yedinennyy institut yedernykh issledovaniy) TITLE: Control of a 680 Nev synchrocyclotron SOURCE: Pribory i tekhnika eksperimenta, no. 3, 1966, 19-22 TOPIC TAGS: synchrocyclotron, particle acceleration, coincidence circuit ABSTRACT: The paper presents a system of control of various synchrocyclotron operating conditions. A phototransducer, having an optico-mechanical connection with a high frequency generator furnishes square pulses of positive polarity. These pulses are used for the regulation of the generator and for synchronizing the operating auxiliary apparatus with the accelerator. A flow chart of this operation is shown. In the continuous mode of operation, the capture and acceleration of the particles occurs in each period of modulation. The synchronization pulses, coincident with the front of the phototransducer pulses, are directed into two channels. In the first of these, the actuating pulses are formed; these pulses move into the exit tube with or without time delay and then into the operator of the high frequency generator. In the second chan-UDC: 621.384.611.2 Card 1/2

| ynchro- oinci- ger ingle of an ol es |
|--------------------------------------|
| l s |
| 'B 1 |
| |
| 7 |
| |
| |
| |
| |
| |
| |
| |
| |
| |

BICH, Ya.A., kand. tekhn. nauk; MURATOV, N.A.; BLISHCHENKO, S.M.; YENDAL'TSEV, B.M.

Rock bumps and efforts to control them in times of the Suchan deposit. Ugol 39 no.5:64-67 My 64. (MIRA 17:8)

1. Vsesoyuznyy nauchno-issledovatel skiy marksheyderskiy institut (for Bich). 2. Shakhta No.21 Suchanskogo mestorozhdeniya (for Muratov). 3. Trest Suchanugol' (for Blishchenko). 4. Shakhta No.10/16 Suchanskogo mestorozhdeniya (for Yendal'tsev).



YENDEN, V. -SHOROKHOV, N.

Facades

Cleaning facades of residential buildings. Zhil.-kom, khoz. 2 no. 7, 1952

MONTHLY LIST OF RUSSIAN ACCESSIONS, LIBRARY OF CONGRESS, NOVEMBER 1952. UNCLASSIFIED.

WENDONOV, Ch.

Budget in the service of the economic and cultural development of the Buryat A.S.S.R. Fin. SSSR 37 no.7:31-34 J1 '63.

(MIRA 16:8)

1. Ministr finansov Buryatskoy ASSR.

(Buryat A.S.S.R.—Budget)

BONDAREV, G.I.; ZINOV'YEV Yo.Sh.; NEPOKLONOV, Yu.A.; YENDOVITSKAYA, I.S.

指数性性性性环境的多数的形式,但有效对现在不可在的结合的过程的特殊的特殊的特殊的一种性心。 他对应的特殊的形式,只是他们的特殊的特别的特殊是不同的的政治的是不可能的

Supply of vitamins C, B₁, B₂ and PP for fish processing workers on fishing craft in the North Atlantic. Vop. pit. 22 no.5:58-60 S-0 '63. (MIRA 17:1)

1. Iz otdela gigiyeny pitaniya (sav. - kand. med. nauk G.I. Bondarev) TSentral'noy nauchno-issledovatel'skoy laboratorii gigiyeny vodnogo transporta, Moskva.

BOHDAREV, G.I.; ZINOV'YEV, Ye.Sh.; NEPOKLOHOV, Yu.A.; YEHDOVITSKAYA, I.S.

Emergy expenditure of fishery workers on travlers fishing in the Barents Sea and North Atlantic. Vop. pit. 21 no.6:40-43 N-D '62. (MIRA 17:5)

1. Iz Tgentral'noy nauchno-issledovatel'skoy laboratorii gigiyeny vodnogo transporta, Moskva.

STANDAY SULFACE STREET STANDARD STANDAR

ARKHANGEL'SKIY, S.N.; YENDOVITSKAYA, T.Y.; NEVEROVICH, Ya.Z.; SOKOLOV, M.V., red.; ALPATOVA, V.V., red.; KOZLOVSKAYA, M.D., tekhn.red.

[Visual aids and experiments for a course in psychology; for pedagogical schools] Nagliadnye posobiia i opyty v kurse psikhologii; dlia pedagogicheskikh uchilishch, Pod red. M.V. Sokolova. Moskva. Gos.uchebno-pedagog. izd-vo M-va prosv. RSFSR, 1958. 103 p. (MIRA 12:1)

(Psychology--Study and teaching)


VENDOVITSKIY, D., metodist

Creative work of the rural vocational schoolteachers. Frof.-tekh.
obr. 22 no.6:5-6 Je '65.

1. Mordovskoye resupblikanskoye upravleniye professional inctekhnicheskogo obrazovaniya.

JEMDÓVITSKIY, V. I.

Budnikov. P.P. and Endovitzkii. V.I. Introducing Birary Kaolins into the Fire-Clay Mix in order to increase its alumina content and refractoriness. Domez, 1932 (1-2) 11-13; Ber. deut. keram. Ges., 13(6) 253-56 (1932.--With the view of improving the quality of some Russian firebrick, experiments were conducted in which binary kaolins were added to the fire-clay mix. Four brick samples, containing 57.68, 56.75, 55.59, and 55.24% SiO₂, respectively, and 36.07, 37.33, 37.99, and 39.25% Al₂O₃, respectively were prepared. These samples were burned at 1350°. The resulting properties were: mechanical strength 239, 255, 289, and 298 kg./sq. cm; refractoriness 1720°, 1735°, 1740°, and 1745°; beginning of deformation under 2 kg./sq. cm. pressure 1395°, 1365° le7t°, and 1400°. A semicommercial plant for the manufacture of brick of the fourth type is recommended.



TENDRIKHOVSKATA, A.T.; PIROZHENIO, A.V.

Fixing automatic feelers on tenter dryer frames. Obm.tekh.
opyt. [MLP] no.10:37-38 '56.
(NIRA 11:11)

(Textile machinery—Attachments)

5.3830 5(4), 5(3)

AUTHORS:

SOY/20-129-4-36/68 Kozlov, P. V., Yendrykhovskaya, A., Kargin, V. A., Academician

TITLE:

Investigation of the Temperature-dependent Transformations in Synthetic Polymers With Rigia Chains

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 129, Nr 4, pp 844-846 (USSR)

ABSTRACT:

The authors investigated polyurea as a typical synthetic polymer with rigid chains. It was produced by polymerization on the boundary of two phases. Phosgene dissolved in benzene was emulsified with hexamethylenediamine dissolved with water. A 7% solution of sodium oleate served as emulsifier. If the 15% hexamethylenediamine solution is saturated with sodium chloride and soda, an amorphous powder with high molecular weight is formed, which is not soluble in any organic solvent with the exception of cresol and formic acid and has a highly ordered structure (Fig 1). The investigation of the temperaturedependent properties was carried out by means of dynamometric scales, a direct dependence of the deformation on temperature being found. Between 230-300°C chemical decomposition already occurs. In order to reduce the temperature at which polyurea is

Card 1/3

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962630010-3"

Investigation of the Temperature-dependent Transformations in Synthetic Polymers With Rigid Chains

transformed, it was plastified according to two methods: a) by the addition of a polymer with clastic chains (copolymer of caprolactam with hexamethylenediamine and adipic acid), b) by swelling in cresol. Figure 1 shows the influence exerted by temperature on polyurea plastified by copolymer. With the addition of 75% copolymer, two processes may be observed: First, transition to the viscous state occurs, after which vitrification follows at a certain temperature. The behavior of polyurea swelled in cresol is shown in figure 3. Also in this case the viscous state occurs with an increase in temperature. On the basis of these results the authors point out the following two possibilities: 1) Hardening of polymers with elastic chains by the admixture of polymers with rigid chains, and 2) reduction of temperature by plastification in order to make working with polymers with rigid chains possible. There are 3 figures and 5 Soviet references.

ASSOCIATION: Card 2/3

Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova (Moscow State University imeni M. V. Lomonosov)

Investigation of the Temperature-dependent Transformations in Synthetic Polymers With Rigid Chains

SUBMITTED:

September 1, 1959

Card 3/3

TITIF. Synthesis and properties of sulfides and phosphides of a remark of the sulfides and phosphides of a remark of the sulfides and phosphides of a remark of the sulfides are sarth metal sulfide, rare metal phosphide synthesis.

ABSTRACT: The sesquisulfides of lanthanum, cerium, prascodymium, and needyntum were obtained by sulfidizing the respective metal nations of a minture of sesquisulfides and oxides with carbon in a vacuum at the sulfides and oxides with

| - T- 12-1 | L 14313-65 ACCESSION NR: AT4047134 Valitum and lantnanum phosp: the respective oxides with prespectively. ASSOCIATION: Institut proble | am material ovedent | Lye AN Ukrssk (Institute | |
|-----------|--|---------------------|--------------------------|--|
| | SUBHITTED: 08Jua64 | ENCT: OA | SUB COUE: 194, 30 | |
| | NO REP SOVE GOS | OTHER; 000 | ATD PRESS: 3136 | |
| • | | | | |

| ACC NR. AP6019225 (A) SOURCE CODE: UR/0073/66/032/002/0115/0118 |
|--|
| AUTHOR: Samsonov, G. V.; Veroykina, L.L.; Yendrzheyevskaya, S. N.; Tikhonova, N.N. |
| ORG: Institute of the Problems of Material Science, AN UkrSSR (Institut Problem materialovedeniya AN UkrSSR) |
| TITLE: Production and some properties of rare-earth phosphides |
| SOURCE: Ukrainskiy khimicheskiy zhurnal, v. 32, no. 2, 1966, 115-118 |
| TOPIC TAGS: rare earth element, phosphide, lanthanum compound, neodymium compound, samarium compound, oxidation |
| ABSTRACT: The literature was reviewed on various methods of producing rare-earth phosphides together with the tabulated data on their crystallochemical proporties (lattice parameters and densities determined from x-ray diffraction patterns). The (lattice parameters and densities determined from x-ray diffraction patterns). The reaction of phosphine (PH3) with rare-earth metals or their oxides was used in this investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidization was carried out investigation for preparing La, Nd, and Sm phosphides. Phosphidizat |

Card 1/2

UDC: 546+661.865

ACC NR: AP6019225

dissolved well in diluted and concentrated HCl and amua regia, and was weakly soluble in H2SO, at any concentration. The NdP was produced either from metallic Nd at 11000 and a 3 hr exposure to an Ar atmosphere, or from Nd2O3 at 13500 and a 3 hr exposure in H. The NdP powder had a black color, a nearly stoichiometric composition, was insoluble in H2O, but dissolved in the same solvents as LaP; SmP of nearly stoichiometric composition was produced from metallic Sm at 9000 after 7 hrs. of phosphidizametric composition was produced from metallic Sm at 9000 after 7 hrs. of phosphidization, and from Sm2O3 at 900-13500 and 2-5 hrs. of phosphidization. From Sm2O3 the SmP tion, and from Sm2O3 at 900-13500. It was in the form of black powder which was formed most efficiently at 1300-13500. It was in the form of black powder which did not change during prolonged storage in air. The SmP dissolved well in HNO3 of varidid not change during prolonged storage in air. The SmP dissolved well in HNO3 of varidid not change during prolonged storage in air. The SmP dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or boiling. Thus, LaP, NdP, and SmP all dissolved well in dilutsolutions either cold or bo

SUB CODE: 07/ SUBM DATE: 30Sep64/ ORIG REF: 007/ OTH REF: 012

Card 2/2

S/057/62/032/008/014/015 B104/b102

AUTHORS: Yendzheyets, G., Molchanov, V. A., Tel'kovskiy, V. G., and Faruk, M. A.

TITLE: Angular distribution of evaporated particles in the irradiation of single crystals with an ion beam

PERIODICAL: Zhurnal tekhnicheskoy fiziki, v. 32, no. 8, 1962, 1032 - 1033

TEXT: The angular distribution of the particles produced when the (100) faces of copper and nickel single crystals were irradiated with argon and neon ions was measured. The diameter of the single crystal surface irradiated was smaller than 8 mm, the distance between target and collector 95 mm. The target temperature was lower than 100°C, the angle of incidence of the ions 20°C. After irradiation five Wehner spots became visible on the collector; four at the corners corresponding to the (110) axis, and one in the center which corresponded to the (100) axis. The density of the spots was determined photometrically. (Fig. 1). The angular distribution of particles and that of the sputtering coefficient do not depend on mass and energy of the ions. There are 3 figures.

Gard 1/2

GUBANOV, M.; YENENKO, B.; BASHKOV, M.; LOBASOV, M.

THE PROPERTY OF COMMENCES AND ADDRESS OF THE PROPERTY OF THE P

Coal and technological progress; our interviews. Sov.shakht. 11 no.2:21-23 F '62. (MIRA 15:1)

1. Direktor Ukrainskogo nauchno-issledovatel skogo instituta Gidrougol (for Gubanov). 2. Direktor Instituta avtomatiki Luganskoy oblasti (for Yenenko). 3. Direktor Luganskogo filiala instituta Giprougleavtomati atsiya (for Bashkov). 4: Direktor Luganskogo filiala Dongiprouglemasha. (for Lobasov).

(Lugansk Province—Goal mines and mining)

NATSH, M. N., inzh.; YENENKO, B. A., inzh.; KOFANOV, G. F., inzh.

Repairing single-model forging and pressing unit in the forge shop after an accident. Mashinostreenie no.5:16-18 S-O '62.

(MERA 16:1)

1. Teplovozostroitel'nyy zaved im. Oktyabr'skey revelyutsii.

(Forging machinery—Maintenance and repair)

YENENKO, G.M., inzh.; STEPANOV, Ye.M., kand. tekhn. nauk; FILIMONOV, Yu.P., kand. tekhn. nauk; KASEN OV, M.A., kand. tekhn. nauk, retsenzent; MAKGVSKIY, G.M., inzh., red.

[Industrial furnaces] Promyshlenrye pechi. Moskva, Mashinostroenie, 1964. 359 p. (MIRA 18:1)

SMIRNOV, B.M., doktor sel'skokhoz. nauk; YENENKO, I.I., aspirant

Controlling offset weeds. Zashch. rast. ot vred. i bol. 9 no.4:
16-17 '64. (MIRA 17:5)

1. Institut sel'skogo khozyaystva Yugo-Vostoka.

AREF YEVA, A.M.; GUTKINA, A.V.; YENENKO, S.O.; MITYUSHIN, V.M.; SHUNGSKAYA, V.Ye.

Cytochemical and electron microscopic study of early changes in ascitic Ehrlich's carcinoma cells following ionizing radiation. Radiobiologia 5 no.3:409-414 '65. (MIRA 18:7)

1. Institut biologicheskoy fiziki AN SSSR, Moskva.

YENENKO, S.O.; SHUNGSKAYA, V.Ye.

Cerebellar cortex cells of adult marmals under supravital conditions. TSitologiia. 6 no.3:380-382 My-Je 164. (MTRA 18:9)

1. Laboratoriya biofiziki zhivykh struktur Instituta biologicheskoy fiziki AN SSSR, Moskva.

AREF'TEVA, A.M.; GUTKINA, A.V.; YENENKO, S.O.; ZARKH, Ye.N.; SHUNGSKAYA,

级。 第一种,我们可以完成了一种的,我们们是这种的数据,就是是我们的,我们就是这种的,我们就是这种的,我们就是这种的,我们就是这种的数据的,我们就是这种的人,就是这个

Cytochemical and luminescence microscopic study of nerve cells in a tissue culture. TSitologiia 7 no.4*511-537 J1-Ag '65.

(MIRA 18:9)

1. Laboratoriya biofiziki zhivykh struktur Instituta biologicheskoy fiziki AN SSSR, Moskva.

YENENKO, O.K.; OZEROV, I.M.; POLOZOV, V.F.; S.PIL'FOGEL', P.V.

Basic properties of the cyclon shale ash of the Central Electric Power station of the "Shale" Combine. Trudy VNIIT no.13:150-161 '64.

(MIRA 18:2)